

# THE ALGEBRAIC CROSSING NUMBER AND THE BRAID INDEX OF KNOTS AND LINKS

Keiko Kawamuro\*

May 3, 2006

## Abstract

It has been conjectured that the algebraic crossing number of link is uniquely determined in minimal braid representation. This conjecture is true for many classes of knots and links.

The Morton-Franks-Williams inequality gives a lower bound for braid index. And sharpness of the inequality on a knot type implies the truth of the conjecture for the knot type.

We prove that there are infinitely many examples of knots and links on which the inequality is not sharp, but the conjecture is still true in these cases. We also show that if the conjecture is true for  $\mathcal{K}$  and  $\mathcal{L}$ , then it is also true for the  $(p, q)$ -cable of  $\mathcal{K}$ , and for the connect sum of  $\mathcal{K}$  and  $\mathcal{L}$ .

## 1 Introduction

The *Braid index* is one of the classical invariants of knots and links. Any knot and link type is presented as a braid closure. The braid index of a link type is the least number of braid strands needed for that.

The *Algebraic crossing number* (or writhe) is a quantity of oriented link diagram counting the crossings with weight  $+1$  (resp.  $-1$ ) for a positive (resp. negative) crossing as shown in the left (resp. middle) sketch of Figure 1. Since it is changed under Reidemeister move I, it is *not* an invariant of link types. However, it has been asked (see [9] page 357 for example):

**Question:** “*Is the algebraic crossing number in a minimal braid representation a link invariant?*”

(Here “minimal” means that the number of braid strands of a link diagram is equal to the braid index of the link type.)

We have known that torus links, closed positive braids with a full twist (including the Lorenz links) [7], 2-bridge links and alternating fibered links [14] and links with braid index  $\leq 3$  [3] have unique algebraic crossing numbers at minimal braid index.

In the following, we overview Section 3 first then go back to Section 2.

Our goal in Section 3 is to approach the question in three ways:

The first way (Theorem 3.5 and its corollaries) is by studying the deficit of the Morton-Franks-Williams (MFW) inequality [12], [7]. It is easy to see that sharpness of the MFW-inequality implies

---

\*The author acknowledges partial support from NSF grants DMS-0405586 and DMS-0306062

the uniqueness of the algebraic crossing number at minimal index. Then how do we answer the question for links on which the inequality is not sharp? In fact we provide infinitely many examples of non-sharp links having unique algebraic crossing numbers at minimal braid index.

The second way is by studying the behavior of the braid index and the algebraic crossing number under the cabling operation. In Theorem 3.10 and Theorem 3.12, we will prove that the uniqueness property is preserved under cabling. Then we have Corollary 3.15 saying “yes” to the question for iterated torus knots.

The third way is by studying the connect sum operation. In Theorem 3.16, we will show that the uniqueness property is preserved under taking the connect sum.

In Section 2, we focus on non-sharpness of the MFW-inequality.

To state the MFW inequality, let  $\mathcal{K}$  be an oriented knot type and let  $K$  be a diagram of  $\mathcal{K}$  on a plane. Focus on one crossing of  $K$  with sign  $\varepsilon$ . Denote  $K_\varepsilon := K$  and let  $K_{-\varepsilon}$  (resp.  $K_0$ ) be the closed braid obtained from  $K_\varepsilon$  by changing the the crossing to the opposite sign  $-\varepsilon$  (resp. resolving the crossing), see Figure 1. The *HOMFLYPT polynomial*  $P_{\mathcal{K}}(v, z) = P_K(v, z)$  satisfies the following




Figure 1: Local views of  $K_+, K_-, K_0$ .

relations (for any choice of a crossing):

$$\begin{aligned} \frac{1}{v}P_{K_+} - vP_{K_-} &= zP_{K_0}. \\ P_{\text{unknot}} &= 1. \end{aligned} \tag{1.1}$$

Now we are ready to state the MFW-inequality.

**Theorem 1.1** [12],[7], *The Morton-Franks-Williams inequality.*

Let  $d_+$  and  $d_-$  be the maximal and minimal degrees of the variable  $v$  of  $P_K(v, z)$ . If a knot type  $\mathcal{K}$  has a closed braid representative  $K$  with braid index  $b_K$  and algebraic crossing number  $c_K$ , then we have

$$c_K - b_K + 1 \leq d_- \leq d_+ \leq c_K + b_K - 1. \tag{1.2}$$

As a corollary,

$$\frac{1}{2}(d_+ - d_-) + 1 \leq b_K, \tag{1.3}$$

giving a lower bound for the braid index  $b_{\mathcal{K}}$  of  $\mathcal{K}$ .

In general, it is hard to determine the braid index. This inequality was the first known result of a general nature relating to the computation of braid index, and it appeared to be quite effective. Jones notes, in [9], that on all but five knots,  $9_{42}, 9_{49}, 10_{132}, 10_{150}, 10_{156}$  in the standard knot table, up to crossing number 10, the MFW inequality is sharp. Furthermore it has been known that the inequality is sharp on all torus links, closed positive  $n$ -braids with a full twist [7], 2-bridge links and fibered alternating links [14].

However, the MFW inequality is not as strong as it appears to be as above. In Theorem 2.5 we give an infinite class of prime links in which the deficit  $D_{\mathcal{K}} := b_{\mathcal{K}} - \frac{1}{2}(d_+ - d_-) - 1$  of the MFW inequality (1.3) can be arbitrarily large. And in Theorem 2.8 we see another infinite class of knots, including  $9_{42}, 9_{49}, 10_{132}, 10_{150}, 10_{156}$ , on which the inequality is not sharp.

Then we may ask “why does non-sharpness occur?” Theorem 2.2 gives a sufficient condition for non-sharpness of the MFW inequality. In fact all the examples in Theorems 2.5 and 2.8 satisfy this sufficient condition.

The idea of Theorem 2.2 is to find knots  $K_{\alpha}$  of known braid index  $= b$  which have a distinguished crossing such that, after changing that crossing to each of the other two possibilities in Figure 1, giving knots or links  $K_{\beta}$  and  $K_{\gamma}$ , it is revealed that  $K_{\beta}$  and  $K_{\gamma}$  each has braid index  $< b$ .

Thanks to Theorem 2.2 one can visually observe the “accumulation” of deficits (for example under the connect sum operation and other linking operation) by looking only at the distinguished crossings which contribute to deficits. See the proof of Theorem 2.5 for details.

**Acknowledgment.** This paper is part of the author’s Ph.D. thesis. She is grateful to her advisor, Joan Birman, for her thoughtful advice and encouragement. She also wishes to thank William Menasco, who told her about the Birman-Menasco diagram, discussed in Section 2.3 and the associated conjecture, when she visited him at SUNY Buffalo. She appreciates many helpful comments by Walter Neumann, Dylan Thurston and Ilya Kofman, and thanks Alexander Stoimenow for showing a preprint and the referee who told her about Stoimenow’s published work. Finally, she especially thanks Mikami Hirasawa, who shared many creative ideas and results about fibered knots including the definition and properties of the enhanced Milnor number.

## 2 Non-sharpness of the Morton-Franks-Williams inequality

### 2.1 Sufficient conditions for non-sharpness

We define the deficit of MFW inequality (Definition 2.1) then give sufficient conditions (Theorem 2.2) for a closed braid on which the inequality is not sharp.

Let  $b_{\mathcal{K}}$  be the braid index of knot type  $\mathcal{K}$ , that is the smallest integer  $b_{\mathcal{K}}$  such that  $\mathcal{K}$  can be represented by a closed  $b_{\mathcal{K}}$ -braid. Let  $b_K, c_K$  denote the braid index and the algebraic crossing number of a braid representative  $K$  of  $\mathcal{K}$ .

**Definition 2.1** *Let*

$$D_{\mathcal{K}} := b_{\mathcal{K}} - \frac{1}{2}(d_+ - d_-) - 1$$

*be the difference of the numbers in (1.3), i.e., of the actual braid index and the lower bound for braid index. Call  $D_{\mathcal{K}}$  the deficit of the MFW inequality for  $\mathcal{K}$ .*

If  $D_{\mathcal{K}} = 0$ , the MFW inequality is sharp on  $\mathcal{K}$ . If  $K$  is a braid representative of  $\mathcal{K}$  let  $D_K^+ := (c_K + b_K - 1) - d_+$  and  $D_K^- := d_- - (c_K - b_K + 1)$ . When  $b_K = b_{\mathcal{K}}$ , we have

$$D_{\mathcal{K}} = \frac{1}{2}(D_K^+ + D_K^-). \tag{2.1}$$

Note that  $D_K^{\pm}$  depends on the choice of braid representative  $K$ , but the deficit  $D_{\mathcal{K}}$  is independent from the choice.

**Theorem 2.2** Assume that  $K$  is a closed braid representative of  $\mathcal{K}$  with  $b_K = b_{\mathcal{K}}$ . Focus on one site of  $K$  and construct  $K_+, K_-, K_0$  (one of the three must be  $K$ ). Let  $\alpha, \beta, \gamma \in \{+, -, 0\}$  and assume that  $\alpha, \beta, \gamma$  are mutually distinct. If  $K_\alpha = K$  and positive destabilization is applicable  $p$ -times to each of  $K_\beta$  and  $K_\gamma$ , then

$$D_K^+ \geq 2p; \quad (2.2)$$

and if  $K_\alpha = K$  and negative destabilization is applicable  $n$ -times to each of  $K_\beta$  and  $K_\gamma$ , then

$$D_K^- \geq 2n. \quad (2.3)$$

Therefore, by (2.1), the MFW inequality is not sharp on  $\mathcal{K}$  if  $p + n > 0$ .

Here is a lemma to prove Theorem 2.2.

**Lemma 2.3** Let  $K$  be a closed braid. Choose one crossing, and construct  $K_+, K_-, K_0$  (one of the three must be  $K$ ). We have

$$d_+(P_{K_+}) \leq \max\{d_+(P_{K_-}) + 2, \quad d_+(P_{K_0}) + 1\} \quad (2.4)$$

$$d_+(P_{K_-}) \leq \max\{d_+(P_{K_+}) - 2, \quad d_+(P_{K_0}) - 1\} \quad (2.5)$$

$$d_+(P_{K_0}) \leq \max\{d_+(P_{K_+}) - 1, \quad d_+(P_{K_-}) + 1\} \quad (2.6)$$

and

$$d_-(P_{K_+}) \geq \min\{d_-(P_{K_-}) + 2, \quad d_-(P_{K_0}) + 1\}$$

$$d_-(P_{K_-}) \geq \min\{d_-(P_{K_+}) - 2, \quad d_-(P_{K_0}) - 1\}$$

$$d_-(P_{K_0}) \geq \min\{d_-(P_{K_+}) - 1, \quad d_-(P_{K_-}) + 1\}.$$

**Proof of Lemma 2.3.** By (1.1), we have  $P_{K_+} = v^2 P_{K_-} + vz P_{K_0}$ . Thus,  $d_+(P_{K_+}) = d_+(v^2 P_{K_-} + vz P_{K_0}) \leq \max\{d_+(v^2 P_{K_-}), d_+(vz P_{K_0})\}$  and we obtain (2.4). The other results follow similarly.  $\square$

Table (2.7) shows the changes of  $c_K, b_K, c_K - b_K + 1$  and  $c_K + b_K - 1$  under stabilization and destabilization of a closed braid.

	$c_K$	$b_K$	$c_K - b_K + 1$	$c_K + b_K - 1$
+ stabilization	+1	+1	0	+2
+ destabilization	-1	-1	0	-2
- stabilization	-1	+1	-2	0
- destabilization	+1	-1	+2	0

(2.7)

Note that  $c_K$  and  $b_K$  are invariant under braid isotopy and exchange moves.

**Proof of Theorem 2.2.** Suppose that  $K = K_\alpha = K_+$ . Suppose we can apply positive destabilization  $k$ -times ( $k \geq p$ ) to  $K_-$ . Let  $\tilde{K}_-$  denote the closed braid obtained after the destabilization. Then we have:

$$\begin{aligned}
d_+(P_{K_-}) + 2 &= d_+(P_{\tilde{K}_-}) + 2 \\
&\leq (c_{\tilde{K}_-} + b_{\tilde{K}_-} - 1) + 2 \\
&= \{(c_{K_-} + b_{K_-} - 1) - 2k\} + 2 \\
&= (c_{K_+} - 2) + b_{K_+} - 1 - 2k + 2 \\
&= (c_{K_+} + b_{K_+} - 1) - 2k = (c_K + b_K - 1) - 2k.
\end{aligned} \quad (2.8)$$

The first equality holds since  $K_-$  and  $\tilde{K}_-$  have the same knot type. The first inequality is the MFW inequality. The second equality follows from Table (2.7).

Similarly, if we can apply positive destabilization  $l$ -times ( $l \geq p$ ) to  $K_0$ , and obtain  $\tilde{K}_0$ , we have

$$\begin{aligned}
 d_+(P_{K_0}) + 1 &= d_+(P_{\tilde{K}_0}) + 1 \\
 &\leq (c_{\tilde{K}_0} + b_{\tilde{K}_0} - 1) + 1 \\
 &= (c_{K_0} + b_{K_0} - 1 - 2l) + 1 \\
 &= (c_{K_+} - 1) + b_{K_+} - 1 - 2l + 1 \\
 &= (c_{K_+} + b_{K_+} - 1) - 2l = (c_K + b_K - 1) - 2l.
 \end{aligned} \tag{2.9}$$

By (2.4), (2.8) and (2.9) we get

$$\begin{aligned}
 d_+(P_K) &= d_+(P_{K_+}) \leq \max\{d_+(P_{K_-}) + 2, \quad d_+(P_{K_0}) + 1\} \\
 &\leq (c_K + b_K - 1) - \min\{2k, 2l\},
 \end{aligned}$$

i.e.,  $D_K^+ \geq \min\{2k, 2l\} \geq 2p$ . When  $K_\alpha = K_-$  or  $K_\alpha = K_0$ , the same arguments work (use (2.5) or (2.6) for these cases in the place of (2.4)) and we get (2.2).

The other inequality (2.3) also holds by the identical argument.  $\square$

## 2.2 Deficit growth

Our goal is to exhibit examples (Theorem 2.5) of prime links on which the deficit of the inequality can be arbitrary large.

**Theorem 2.4** *Knot type  $\mathcal{K} = 9_{42}$  has a braid representative  $K = K_+$  (see Figure 2) satisfying the sufficient condition in Theorem 2.2.*




Figure 2: Knot  $9_{42}$  satisfies the conditions of Theorem 2.2

**Proof of Theorem 2.4.** It is known that  $9_{42}$  has braid index = 4 and deficit  $D_{9_{42}} = 1$ . Let  $K = K_+$  be its braid representative of the minimal braid index as in Figure 2. Construct  $K_-, K_0$  by changing the shaded crossing. Sketches show that both  $K_-, K_0$  can be positively destabilized. Thus by Theorem 2.2,  $D_K^+ \geq 2$  and  $D_{9_{42}} \geq 1$ .  $\square$

**Theorem 2.5** *For any positive integer  $n$ , there exists a prime link  $L$  whose deficit  $D_L \geq n$ .*

**Proof of Theorem 2.5.** We prove the theorem by exhibiting examples. For  $n \in \mathbb{N}$  let  $\mathcal{A}^n(9_{42})$  be the closure of  $n$ -copies of  $9_{42}$  linked each other by two full twists as in the left sketch of Figure 3. Since the braid index  $b_{9_{42}} = 4$  and  $\mathcal{A}^n(9_{42})$  is an  $n$ -component link, we know the braid index of




Figure 3: Prime link  $\mathcal{A}^5(9_{42})$  and 2-component link  $\mathcal{A}$ .

$\mathcal{A}^n(9_{42})$  is  $4n$ . This construction gives a braid representative with  $4n$ -strands and  $n$  distinguished (shaded in the left sketch) crossings.

In the following we will see that each of the shaded crossing contributes to the deficit.

Let  $\mathcal{K} := \mathcal{A}^2(9_{42})$  and let  $K$  be the braid representative of  $\mathcal{K}$  as in Figure 3. Let  $K_{--}, K_{-0}, K_{0-}, K_{00}$  be the links obtained from  $K$  by changing the two shaded crossings. We repeat the discussion of the proof of Theorem 2.2: We have

$$\begin{aligned}
d_+(P_{K_{--}}) + (2 + 2) &= d_+(P_{\tilde{K}_{--}}) + 4 \\
&\leq (c_{\tilde{K}_{--}} + b_{\tilde{K}_{--}} - 1) + 4 \\
&= \{(c_{K_{--}} + b_{K_{--}} - 1) - 2 \cdot 2\} + 4 \\
&= (c_K - 4) + b_K - 1 - 2 \cdot 2 + 4 \\
&= (c_K + b_K - 1) - 2 \cdot 2.
\end{aligned}$$

Similarly,

$$\begin{aligned}
d_+(P_{K_{-0}}) + (2 + 1) &\leq (c_K + b_K - 1) - 2 \cdot 2, \\
d_+(P_{K_{0-}}) + (1 + 2) &\leq (c_K + b_K - 1) - 2 \cdot 2, \\
d_+(P_{K_{00}}) + (1 + 1) &\leq (c_K + b_K - 1) - 2 \cdot 2.
\end{aligned}$$

Thus,

$$\begin{aligned}
d_+(P_K) &= \max\{d_+(P_{K_{--}}) + 4, d_+(P_{K_{-0}}) + 3, d_+(P_{K_{0-}}) + 3, d_+(P_{K_{00}}) + 2\} \\
&\leq (c_K + b_K - 1) - 2 \cdot 2
\end{aligned}$$

and  $D_{\mathcal{K}} \geq \frac{1}{2}D_K^+ \geq \frac{1}{2}(2 \cdot 2) = 2$ .

Similar arguments work when  $\mathcal{K} = \mathcal{A}^n(9_{42})$  for  $n \geq 3$  and we have  $D_{\mathcal{A}^n(9_{42})} \geq \frac{1}{2}D_{\mathcal{A}^n(9_{42})}^+ \geq \frac{1}{2}(2 \cdot n) \geq n$ .

The 2-component link  $\mathcal{A}$  of the right sketch is hyperbolic [15]. Pair  $(S^3, \mathcal{A}^n(9_{42}) \cup z\text{-axis})$  is an  $n$ -fold cover of  $(S^3, \mathcal{A})$  branched at  $z$ -axis. Therefore, by [17] we can conclude that  $\mathcal{A}^n(9_{42})$ 's are all prime except for finitely many  $n$ 's.  $\square$

**Remark 2.6** By taking the connected sum of knots on which the MFW inequality is non-sharp, one can also construct examples of (non-prime) knots with arbitrarily large deficits. This fact follows not only by Theorem 2.2 but also by the definition of HOMFLYPT polynomial (1.1) and the additivity of braid indices under connected sums [4].

### 2.3 Birman-Menasco diagram

As an application of Theorem 2.2, we study another infinite class of knots including all the Jones' five knots  $(9_{42}, 9_{49}, 10_{132}, 10_{150}, 10_{156})$  on which the MFW inequality is not sharp. We call the block-strand diagram (see [6] for definition) of Figure 4 the Birman-Menasco (BM) block-strand diagram.




Figure 4: The Birman-Menasco diagram  $BM_{x,y,z,w}$ .

**Definition 2.7** Let  $BM_{x,y,z,w}$ , where  $x, y, z, w \in \mathbb{Z}$ , be the knot (or the link) type which is obtained by assigning  $x$  positive half-twists (resp.  $y, z, w$ ) to the braid block  $X$  (resp.  $Y, Z, W$ ) of the BM diagram.

Recall that on all but only five knots  $(9_{42}, 9_{49}, 10_{132}, 10_{150}, 10_{156})$  up to crossing number 10 the MFW inequality is sharp. An interesting property of the BM diagram is that it carries all the five knots. Namely, we have  $9_{42} = BM_{-1,1,-2,-1} = BM_{-1,-2,-2,2}$ ,  $9_{49} = BM_{-1,1,1,2}$ ,  $10_{132} = BM_{-1,-2,-2,-2}$ ,  $10_{150} = BM_{3,-2,-2,2} = BM_{-1,2,-2,2} = BM_{-1,-2,2,2} = BM_{-1,1,2,-1} = BM_{3,1,-2,-1}$ , and  $10_{156} = BM_{-1,1,1,-2}$ .

We have the following theorem, which was conjectured informally by Birman and Menasco:

**Theorem 2.8** There are infinitely many  $(x, y, z, w)$ 's such that the MFW inequality is not sharp on  $BM_{x,y,z,w}$ .

**Lemma 2.9** We have  $D_{BM_{x,y,z,w}}^+ \geq 2$ .

**Proof of Lemma 2.9.** Change the BM diagram into the diagram in sketch (1) of Figure 5 by braid isotopy and denote it by  $K$ . Focus on the crossing shaded in sketch (1). Regard  $K = K_-$ . We can apply positive destabilization once to  $K_+$  and obtain the diagram in sketch (2-2). We also




Figure 5: BM-diagram satisfies the sufficient condition

can apply positive destabilization once to  $K_0$  as we can see in the passage sketch (3-1)  $\Rightarrow$  (3-2)  $\Rightarrow$  (3-3). Therefore by Theorem 2.2 we have  $D_{BM_{x,y,z,w}}^+ \geq 2$  for any  $(x, y, z, w)$ . □

It remains to prove that there exist infinitely many  $(x, y, z, w)$ 's such that the braid index of  $BM_{x,y,z,w}$  is 4. We introduce  $\mathcal{K}_n := BM_{-1,-2,n,2}$  and will show that for all  $m \geq 1$  the braid index of  $\mathcal{K}_{2m}$  is 4. (Note that  $\mathcal{K}_{-2} = 9_{42}, \mathcal{K}_2 = 10_{150}$  and  $\mathcal{K}_{2m}$  is a knot.) It will then follow, thanks to Lemma 2.9, that the MFW inequality cannot be sharp on any  $\mathcal{K}_{2m}$ ,  $m \geq 1$ .

In order to do this, we use the *enhanced Milnor number*  $\lambda$  defined by Neumann and Rudolph [16]. Recall that the fiber surface of a fiber knot is obtained by plumbing and unplumbing Hopf bands [8]. This  $\lambda$  is an invariant of fibered knots and links counting algebraically the number of negative Hopf bands to get the fiber surface.

**Lemma 2.10** *All  $\mathcal{K}_n$  ( $n \geq 2$ ) are fibered and have the enhanced Milnor number  $\lambda = 1$ .*

**Proof of Lemma 2.10.** Sketch (1) of Figure 6 is the standard Bennequin surface of  $\mathcal{K}_n$ . We compress it twice as in the passage sketch (1)  $\Rightarrow$  (2) along the disks bounded by dotted circles in sketch (1). Next, deplumb positive Hopf bands as much as possible as in the passage sketch (2)  $\Rightarrow$  (3)  $\Rightarrow$  (4) = (5). Then isotope the surface until we get sketch (8). These operations do not change the enhanced Milnor number.

We apply Melvin and Morton's trick [11] p.167, as in the passage sketch (8)  $\Rightarrow$  (9). We remark that the enhanced Milnor number is invariant under this trick.

The surface of sketch (9) = (10), whose boundary is Pretzel link  $P(-2, 0, 2)$ , is plumbing of a positive Hopf band and a negative Hopf band. Thus it has  $\lambda = 1$  so does  $\mathcal{K}_n$ . □

Here we summarize Xu's classification of 3-braids [21]. Let  $\sigma_1, \sigma_2$  be the standard generators of  $B_3$  the braid group of 3-strings satisfying  $\sigma_1\sigma_2\sigma_1 = \sigma_2\sigma_1\sigma_2$ . Let  $a_1 := \sigma_1, a_2 := \sigma_2$  and  $a_3 := \sigma_2\sigma_1\sigma_2^{-1}$ . We can identify them with the twisted bands in Figure 7.

Let  $\alpha := a_1a_3 = a_2a_1 = a_3a_2$ . If  $w \in B_3$  let  $\bar{w}$  denote  $w^{-1}$ .

**Theorem 2.11** (Xu [21].) *Every conjugacy class in  $B_3$  can be represented by a shortest word in  $a_1, a_2, a_3$  uniquely up to symmetry. And the word has one of the three forms:*

$$(1)\alpha^k P, \quad (2)N\bar{\alpha}^k, \quad (3)NP.$$

where  $k \geq 0$  and  $\bar{N}, P$  are positive words and the arrays of subscripts of the words are non-decreasing.

The next is another lemma for Theorem 2.8:

**Lemma 2.12** *If a closed 3-braid has  $\lambda = 1$  and is a knot, then up to symmetry it has one of the following forms:*

$$\begin{aligned} A_x &:= \bar{a}_3 \bar{a}_2 (a_1)^x, \quad x \geq 2, \text{ even}, \\ B_{x,y} &:= \bar{a}_3 \bar{a}_3 (a_1)^x (a_2)^y, \quad x, y \geq 3, \text{ odd}, \\ C_{x,y,z} &:= \bar{a}_2 (a_1)^x (a_2)^y (a_3)^z, \quad x+z = \text{odd}, \quad y = \text{even}, \quad x, y, z \geq 1, \\ D_{x,y,z,w} &:= \bar{a}_2 (a_1)^x (a_2)^y (a_3)^z (a_1)^w, \quad x, y \geq 2, \quad z, w \geq 1. \end{aligned}$$




Figure 6: Deformation of  $\mathcal{K}_n$




Figure 7: Xu's band generators

**Proof of Lemma 2.12.** Assume we have a word  $w \in B_3$ . By Theorem 2.11,  $w$  has one of the following forms:

Case (1)-1	$w = \alpha^k$	$k \geq 1$
Case (1)-2	$w = \alpha^k P$	$k \geq 1$
Case (1)-3	$w = P$	no $\alpha$ part
Case (2)-1	$w = \bar{\alpha}^k$	$k \geq 1$
Case (2)-2	$w = N\bar{\alpha}^k$	$k \geq 1$
Case (2)-2	$w = N$	no $\bar{\alpha}$ part
Case (3)	$w = NP$	18 cases to study

In this proof, we use the simplified notations:

symbol	meaning	change in $\lambda$
$i$	$a_i$ for $i = 1, 2, 3$ .	—
$=$	same conjugacy class	0
$\longrightarrow$	deplumb positive-Hopf bands	0
$\implies$	deplumb negative-Hopf bands	$\geq 1$
$\approx$	Melvin-Morton trick [11]	0
$\rightsquigarrow$	composition of deplumbings of $\pm$ Hopf bands	$\geq 0$

These are formulae we use:

$$\alpha^2 \longrightarrow \alpha, \text{ Figure 8} \quad (2.10)$$

$$\alpha 123 \longrightarrow \alpha, \text{ Figure 9} \quad (2.11)$$

$$\bar{i}(i-1)i \approx \bar{i} \bar{i-1}i, \text{ Melvin-Morton trick} \quad (2.12)$$

$$i(i+1)\bar{i} \approx i \overline{i+1} \bar{i}, \text{ Melvin-Morton trick} \quad (2.13)$$




Figure 8:  $\alpha^2 \longrightarrow \alpha$ .




Figure 9:  $\alpha 123 \longrightarrow \alpha$ .

Now we study each case.

**Case (1)-1.** By (2.10), we have  $w = \alpha^k \rightarrow \alpha (= \text{unknot})$ . Thus  $w$  has  $\lambda = 0$ .

**Case (1)-2.** By (2.10) up to permutation of  $\{1, 2, 3\}$  we have  $\alpha^k P \rightarrow \alpha P \rightarrow \alpha(123123 \cdots \cdots)$ . Thanks to (2.11) we have

$$\alpha \overbrace{123123 \cdots \cdots}^{\text{length} = l} \rightarrow \alpha \overbrace{123123 \cdots \cdots}^{\text{length} = l - 3} \quad \text{for } l \geq 3.$$

If  $l = 1, 2$ , we have  $\alpha 1 = 211 \rightarrow \alpha$  and  $\alpha 12 = 2112 \rightarrow \alpha$ . Thus  $w$  has  $\lambda = 0$ .

**Case (1)-3.** Assume  $w = P$ . There are three possible cases:

$$P \rightarrow (123)^n, \quad P \rightarrow (123)^n 1 \quad \text{and} \quad P \rightarrow (123)^n 12 \quad \text{where } n \geq 0.$$

If  $w$  satisfies the first case, it is proved that  $(123)^n$  is not fibered in Theorem 3.2 of [19], where Stoimenow determines fibreeness of strongly quasi-positive 3-braid links. Therefore,  $w$  is not fibered.

The second case can be reduced to the first case, since  $(123)^n 1 = 1(123)^n \rightarrow (123)^n$ .

For the third case, since  $(123)^n 12 = 2(123)^n 1 = \alpha(231)^n \rightarrow \alpha$ ,  $w$  has  $\lambda = 0$ .

**Case (2)-1.** By (2.10),  $w = \bar{\alpha}^k \Rightarrow \bar{\alpha}$  and  $w$  has  $\lambda = 2(k - 1) \neq 1$ .

**Case (2)-2.** Suppose  $w = N\bar{\alpha}^k$  where  $k \geq 1$ .

If  $w = \bar{i} \bar{\alpha}$ , we have  $\bar{i} \bar{\alpha} \Rightarrow \bar{\alpha}$  and  $w$  has  $\lambda = 1$ . However, the closure of  $w$  has more than one component and it does not satisfy the condition of the lemma.

If  $w \neq \bar{i} \bar{\alpha}$ , we have  $N\bar{\alpha}^k \Rightarrow \bar{\alpha}$  by (2.11), and  $w$  has  $\lambda \geq 2$ .

**Case (2)-3.** Suppose  $w = N$ . There are three possible cases:

$$N \Rightarrow (\bar{3} \bar{2} \bar{1})^n, \quad N \Rightarrow (\bar{3} \bar{2} \bar{1})^n \bar{3} \quad \text{and} \quad N \Rightarrow (\bar{3} \bar{2} \bar{1})^n \bar{3} \bar{2} \quad \text{where } n \geq 0.$$

For the first case,  $w$  is not fibered [19].

For the second case, if  $n = 0$  then  $w$  has  $\lambda = 1$  if and only if  $w = \bar{3} \bar{3}$ . However this has two components. If  $n \geq 1$ , since  $(\bar{3} \bar{2} \bar{1})^n \bar{3} \Rightarrow (\bar{3} \bar{2} \bar{1})^n$  it can be reduced to the first case.

For the third case, if  $n = 0$  then  $w$  has  $\lambda = 1$  if and only if  $w = \bar{3} \bar{3} \bar{2}$ . However it has two components. If  $n \geq 1$ , we have  $(\bar{3} \bar{2} \bar{1})^n \bar{3} \bar{2} = \bar{2} \bar{3} (\bar{2} \bar{1} \bar{3})^n = \bar{\alpha} (\bar{2} \bar{1} \bar{3})^n \Rightarrow \bar{\alpha}$  and  $w$  has  $\lambda \geq 3n$ .

**Case (3).** Assume  $w = NP$ . Let  $w'$  be a word obtained from  $w$  by deplumbing  $\pm$  Hopf bands sufficiently enough times, i.e.,  $w \rightsquigarrow w'$ . This  $w'$  has one of the following 18 forms up to permutation of  $\{1, 2, 3\}$ .

case	$w'$
i	$(\bar{2} \bar{1} \bar{3})^k (123)^l \quad k, l \geq 1$
ii	$(\bar{2} \bar{1} \bar{3})^k (123)^l 1 \quad k \geq 1, l \geq 0$
iii	$(\bar{2} \bar{1} \bar{3})^k (123)^l 12 \quad k \geq 1, l \geq 0 \quad \text{not shortest word}$
iv	$\bar{3} (\bar{2} \bar{1} \bar{3})^k (123)^l \quad k \geq 0, l \geq 1 \quad \text{not shortest word}$
v	$\bar{3} (\bar{2} \bar{1} \bar{3})^k (123)^l 1 \quad k, l \geq 0$
vi	$\bar{3} (\bar{2} \bar{1} \bar{3})^k (123)^l 12 \quad k, l \geq 0$
vii	$\bar{1} \bar{3} (\bar{2} \bar{1} \bar{3})^k (123)^l \quad k \geq 0, l \geq 1$
viii	$\bar{1} \bar{3} (\bar{2} \bar{1} \bar{3})^k (123)^l 1 \quad k, l \geq 0 \quad \text{not shortest word}$
ix	$\bar{1} \bar{3} (\bar{2} \bar{1} \bar{3})^k (123)^l 12 \quad k, l \geq 0$

i'	$(\bar{1} \bar{3} \bar{2})^k (123)^l$	$k, l \geq 1$
ii'	$(\bar{1} \bar{3} \bar{2})^k (123)^l 1$	$k \geq 1, l \geq 0$ not shortest word
iii'	$(\bar{1} \bar{3} \bar{2})^k (123)^l 12$	$k \geq 1, l \geq 0$
iv'	$\bar{2}(\bar{1} \bar{3} \bar{2})^k (123)^l$	$k \geq 0, l \geq 1$
v'	$\bar{2}(\bar{1} \bar{3} \bar{2})^k (123)^l 1$	$k, l \geq 0$
vi'	$\bar{2}(\bar{1} \bar{3} \bar{2})^k (123)^l 12$	$k, l \geq 0$ not shortest word
vii'	$\bar{3} \bar{2}(\bar{1} \bar{3} \bar{2})^k (123)^l$	$k \geq 0, l \geq 1$ not shortest word
viii'	$\bar{3} \bar{2}(\bar{1} \bar{3} \bar{2})^k (123)^l 1$	$k, l \geq 0$
ix'	$\bar{3} \bar{2}(\bar{1} \bar{3} \bar{2})^k (123)^l 12$	$k, l \geq 0$

Since words of case iii, iv, viii, ii', vi' and vii' are not shortest (reducible) we eliminate them from the list.

These are reduction formulae we use:

$$(\bar{2} \bar{1} \bar{3})(123) \xrightarrow{(2.19)} \bar{2}(3\bar{2}) = \bar{1}\bar{2} \bar{2} \implies \bar{1}\bar{2} \quad (2.14)$$

$$(\bar{1}\bar{2})(123) \xrightarrow{(2.28)} \bar{1}\bar{1}\bar{2} \longrightarrow \bar{1}\bar{2} \quad (2.15)$$

$$(\bar{2} \bar{1} \bar{3})(\bar{1}\bar{2}) \xrightarrow{(2.17)} \bar{1}\bar{3} \bar{2} = \bar{2} \bar{2}\bar{3} \implies = \bar{1}\bar{2} \quad (2.16)$$

$$(\bar{2} \bar{1} \bar{3})1 \approx \bar{2} \bar{1}31 = \bar{1}\bar{2} \bar{2}1 \implies = \bar{1}\bar{1}\bar{3} \longrightarrow \bar{1}\bar{3} \quad (2.17)$$

$$(\bar{2} \bar{1} \bar{3})\bar{1}\bar{3} \xrightarrow{(2.17)} \bar{1}\bar{3} \bar{3} \implies \bar{1}\bar{3} \quad (2.18)$$

$$\bar{1} \bar{3}(123) = \bar{1}\bar{2}\bar{2}\bar{1}\bar{3} \longrightarrow \bar{1}\bar{2}\bar{1}\bar{3} = \bar{3}\bar{1} \bar{1}\bar{3} \implies = \bar{3}\bar{3}\bar{2} \longrightarrow \bar{3}\bar{2} \quad (2.19)$$

$$3\bar{2}(123) \xrightarrow{(2.28)} 3(\bar{1}\bar{2}) = \bar{1}\bar{3}\bar{3} \longrightarrow \bar{1}\bar{3} = \bar{3}\bar{2} \quad (2.20)$$

$$\begin{aligned} (123)\bar{1}\bar{3} &\approx 123\bar{1} \bar{3} = \bar{1}\bar{3}\bar{2}\bar{2}\bar{3} \\ &\longrightarrow \bar{1}\bar{3}\bar{2}\bar{3} = \bar{1}\bar{3} \bar{3}\bar{1} \implies \bar{1}\bar{3}\bar{1} \longrightarrow \bar{1}\bar{3} \end{aligned} \quad (2.21)$$

$$12\bar{1} \bar{3} = \bar{1}\bar{3} \bar{3}\bar{1} \implies \bar{1}\bar{3}\bar{1} \longrightarrow \bar{2}\bar{1} = \bar{1}\bar{3} \quad (2.22)$$

$$(\bar{1} \bar{3} \bar{2})(123) \xrightarrow{(2.26)} \bar{3}\bar{1}\bar{3} = \bar{3}\bar{3}\bar{2} \implies \bar{3}\bar{2} = \bar{1}\bar{3} \quad (2.23)$$

$$(\bar{1} \bar{3} \bar{2})\bar{1}\bar{3} = \bar{1} \bar{3} \bar{1}\bar{2} \bar{2} \implies \approx \bar{1}\bar{3}\bar{1}\bar{2} = \bar{1} \bar{1}\bar{3}\bar{3} \longrightarrow \implies \bar{1}\bar{3} \quad (2.24)$$

$$\bar{1}\bar{3}(123) \approx \bar{1} \bar{3}\bar{1}\bar{2}\bar{3} = \bar{1}\bar{2}\bar{2}\bar{1}\bar{3} \longrightarrow \implies \bar{3}\bar{1}\bar{3} \longrightarrow \bar{3}\bar{2} = \bar{1}\bar{3} \quad (2.25)$$

$$(\bar{1} \bar{3} \bar{2})12 = \bar{1}\bar{2}\bar{3} \bar{3}\bar{2} \implies = \bar{1}\bar{2}\bar{2}\bar{1} \longrightarrow \bar{1}\bar{2}\bar{1} = \bar{3}\bar{1} \bar{1} \implies \bar{3}\bar{1} \quad (2.26)$$

$$(\bar{1} \bar{3} \bar{2})\bar{3}\bar{1} \xrightarrow{(2.16)} \bar{3}\bar{1} \text{ permutation of (2.16)} \quad (2.27)$$

$$\bar{2}(123) \approx \bar{2} \bar{1}\bar{2}\bar{3} = \bar{2}\bar{3}\bar{3}\bar{2} \longrightarrow \bar{2}\bar{3}\bar{2} = \bar{1}\bar{2} \bar{2} \implies \bar{1}\bar{2} \quad (2.28)$$

$$\bar{1}\bar{2} (\bar{1} \bar{3} \bar{2}) \approx \bar{1}\bar{2}\bar{1} \bar{3} \bar{2} = \bar{1}\bar{3} \bar{3}\bar{1}\bar{2} \implies = \bar{2}\bar{1}\bar{1}\bar{2} \longrightarrow \bar{2}\bar{1}\bar{2} \implies \bar{1}\bar{2} \quad (2.29)$$

**Sublemma 2.13** For  $k, l \geq 1$ , we have:

$$(\bar{2} \bar{1} \bar{3})^k (123)^l \rightsquigarrow \bar{1}\bar{2} \quad (2.30)$$

$$(\bar{1} \bar{3} \bar{2})^k (123)^l \rightsquigarrow \bar{1}\bar{3} \quad (2.31)$$

Either case, the increase of  $\lambda$  is  $\geq 2$ .

**Proof.** From (2.14), (2.16) and (2.17) we obtain (2.30). Similarly, (2.31) follows from (2.23), (2.24) and (2.25).  $\square$

**Case 3-i.** By (2.30), our  $w$  is fibered and  $\lambda \geq 2$ .

**Case 3-ii.** If  $k \geq 1, l = 0$ ,

$$w \rightsquigarrow w' = (\bar{2} \bar{1} \bar{3})^k 1 \xrightarrow{(2.17)} (\bar{2} \bar{1} \bar{3})^{k-1} \bar{1} \bar{3} \xrightarrow{(2.18)} \dots \xrightarrow{(2.18)} \bar{1} \bar{3}$$

and  $w$  has  $\lambda = 1$  if and only if  $w = \bar{2} \bar{1} \bar{3} 1^x$  for some  $x \geq 1$ . However it has 2 or 3 components and it does not satisfy the condition of Lemma 2.12. If  $k, l \geq 1$ ,

$$w \rightsquigarrow w' = (\bar{2} \bar{1} \bar{3})^k (123)^l 1 \xrightarrow{(2.30)} (\bar{1} \bar{2}) 1 \longrightarrow \bar{1} \bar{2}$$

and  $w$  has  $\lambda \geq 2$ .

**Case 3-v.** When  $k = l = 0$ ,  $w$  has  $\lambda = 1$  if and only if  $w = \bar{3} \bar{3} 1^x$  for some  $x \geq 1$ , which has more than 1 component. If  $k \geq 1$  and  $l = 0$ ,

$$w \rightsquigarrow w' = \bar{3} (\bar{2} \bar{1} \bar{3})^k 1 \xrightarrow{(2.18)} \dots \xrightarrow{(2.18)} \bar{1} \bar{3}$$

and  $\lambda \geq 2$ . If  $k = 0, l \geq 1$ ,

$$w \rightsquigarrow w' = \bar{3} (123)^l 1 = (123)^l \bar{1} \bar{3} \xrightarrow{(2.21)} \dots \xrightarrow{(2.21)} \bar{1} \bar{3}.$$

Thus  $w$  has  $\lambda = 1$  if and only if  $w = \bar{3} 1^x 2^y 3^z 1^w$  for  $x, y, z, w \geq 1$ . If  $k, l \geq 1$ ,

$$w \rightsquigarrow w' = \bar{3} (\bar{2} \bar{1} \bar{3})^k (123)^l 1 \xrightarrow{(2.30)} \bar{3} (\bar{1} \bar{2}) 1 = \bar{3} \bar{1} \bar{1} \bar{3} \longrightarrow \implies \bar{1} \bar{3}$$

and  $\lambda \geq 2$ .

**Case 3-vi.** If  $k = l = 0$ ,

$$w \rightsquigarrow w' = \bar{3} 12 = 2\bar{2}\bar{1} \longrightarrow 2\bar{1}.$$

Therefore,  $w$  has  $\lambda = 1$  if and only if  $w = \bar{3} \bar{3} 1^x 2^y$  for  $x, y \geq 1$ . If  $k = 0, l \geq 1$

$$w \rightsquigarrow w' = \bar{3} (123)^l 12 = (123)^l 12\bar{3} \longrightarrow (123)^l \bar{1} \bar{3} \xrightarrow{(2.21)} \dots \xrightarrow{(2.21)} \bar{1} \bar{3}$$

and  $w$  has  $\lambda = 1$  if and only if  $w = \bar{3} 1^x 2^y 3^z 1^w 2^v$  for some  $x, y, z, w, v \geq 1$ . When  $k \geq 1, l = 0$ ,

$$w \rightsquigarrow w' = \bar{3} (\bar{2} \bar{1} \bar{3})^k 12 \longrightarrow (\bar{2} \bar{1} \bar{3})^k \bar{1} \bar{3} \xrightarrow{(2.18)} \dots \xrightarrow{(2.18)} \bar{1} \bar{3}$$

and  $\lambda \geq 2$ . If  $k, l \geq 1$ ,

$$w \rightsquigarrow w' \xrightarrow{(2.30)} \bar{3} (\bar{1} \bar{2}) 12 = \bar{3} \bar{1} \bar{1} \bar{3} 2 \longrightarrow \bar{3} \bar{1} \bar{3} 2 = 2\bar{3} \bar{3} 2 \longrightarrow \implies 2\bar{3}$$

and  $\lambda \geq 2$ .

**Case 3-vii.** If  $k = 0, l \geq 1$ ,

$$w \rightsquigarrow w' = \bar{1} \bar{3} (123)^l \xrightarrow{(2.19)} \bar{3} \bar{2} (123)^{l-1} \xrightarrow{(2.20)} \dots \xrightarrow{(2.19)} \bar{3} \bar{2}$$

and  $\lambda = 1$  if and only if  $w = \bar{1} \bar{3} 1^x 2^y 3^z$  for some  $x, y, z \geq 1$ . If  $k, l \geq 1$ ,

$$w \rightsquigarrow w' \xrightarrow{(2.30)} \bar{1} \bar{3} (\bar{1} \bar{2}) = \bar{2} \bar{1} \bar{3} 1 \xrightarrow{(2.17)} \bar{1} \bar{3}$$

and  $\lambda \geq 2$ .

**Case 3-ix.** When  $k = l = 0$ ,

$$w \rightsquigarrow w' = \bar{1} \bar{3} 12 = \bar{1}22\bar{1} \longrightarrow \Longrightarrow \bar{1}2.$$

Thus  $w$  has  $\lambda = 1$  if and only if  $w = \bar{1} \bar{3} 1^x 2^y$  for some  $x, y \geq 1$ . When  $k = 0, l \geq 1$

$$w \rightsquigarrow w' = \bar{1} \bar{3}(123)^l 12 \xrightarrow{(2.19)} \xrightarrow{(2.20)} \bar{1}312 = 3\bar{2}12 \approx 3\bar{2} \bar{1}2 = \bar{1}33\bar{1} \longrightarrow \Longrightarrow \bar{1}3$$

and  $\lambda \geq 2$ . When  $k \geq 1, l = 0$

$$w \rightsquigarrow w' = \bar{1} \bar{3}(\bar{2} \bar{1} \bar{3})^k 12 = 12\bar{1} \bar{3}(\bar{2} \bar{1} \bar{3})^k \xrightarrow{(2.22)} \bar{1}\bar{3}(\bar{2} \bar{1} \bar{3})^k \xrightarrow{(2.18)} \dots \xrightarrow{(2.18)} \bar{1}\bar{3}$$

and  $\lambda \geq 2$ . When  $k, l \geq 1$

$$w \rightsquigarrow w' \xrightarrow{(2.30)} \bar{1} \bar{3}(\bar{1}\bar{2})12 = 12\bar{1} \bar{3}1\bar{2} \xrightarrow{(2.22)} \bar{1}\bar{3}1\bar{2} = \bar{2}11\bar{2} \longrightarrow \Longrightarrow \bar{1}\bar{2}$$

and  $\lambda \geq 2$ .

**Case 3-i'.** By (2.31) our  $w$  is fibered and  $\lambda \geq 2$ .

**Case 3-iii'.** When  $k > l = 0$ ,

$$w \rightsquigarrow w' = (\bar{1} \bar{3} \bar{2})^k 12 \xrightarrow{(2.26)} (\bar{1} \bar{3} \bar{2})^{k-1} 2\bar{1} \xrightarrow{(2.27)} \dots \xrightarrow{(2.27)} 3\bar{1}$$

and  $\lambda \geq 2$ . When  $k, l \geq 1$ ,

$$w \rightsquigarrow w' \xrightarrow{(2.31)} (\bar{1}\bar{3})12 \approx \bar{1} \bar{3} 12 = \bar{1}22\bar{1} \longrightarrow \Longrightarrow 3\bar{1}$$

and  $\lambda \geq 2$ .

**Case 3-iv'.** When  $k = 0, l \geq 1$ ,

$$w \rightsquigarrow w' = \bar{2}(123)^l \xrightarrow{(2.28)} \bar{1}\bar{2}(123)^{l-1} \xrightarrow{(2.15)} \dots \xrightarrow{(2.15)} \bar{1}\bar{2}.$$

Thus  $w$  has  $\lambda = 1$  if and only if  $w = \bar{2}1^x 2^y 3^z$  for  $x, y, z \geq 1$ . To make the braid closure one component, we further require  $x + z = \text{odd}$ . If  $k, l \geq 1$ ,

$$w \rightsquigarrow w' \xrightarrow{(2.31)} \bar{2}(\bar{1}\bar{3}) = \bar{1}\bar{2} \bar{2} \Longrightarrow \bar{1}\bar{2}$$

and  $\lambda \geq 2$ .

**Case 3-v'.** When  $k = l = 0$ ,

$$w \rightsquigarrow w' = \bar{2}1.$$

Thus  $w$  has  $\lambda = 1$  if and only if  $w = \bar{2} \bar{2}1^x$ , which has 2 or 3 components. When  $k = 0$  and  $l \geq 1$ ,

$$w \rightsquigarrow w' = \bar{2}(123)^l 1 \xrightarrow{(2.28)} \bar{1}\bar{2}(123)^{l-1} 1 \xrightarrow{(2.15)} \dots \xrightarrow{(2.15)} \bar{1}\bar{2}1 = 11\bar{2} \longrightarrow \bar{1}\bar{2}$$

thus  $w$  has  $\lambda = 1$  if and only if  $w = \bar{2} 1^x 2^y 3^z 1^w$  for some  $x, y, z, w \geq 1$ . When  $k \geq 1, l = 0$ ,

$$w \rightsquigarrow w' = \bar{2}(\bar{1} \bar{3} \bar{2})^k 1 = \bar{1}\bar{2}(\bar{1} \bar{3} \bar{2})^k \xrightarrow{(2.29)} \dots \xrightarrow{(2.29)} \bar{1}\bar{2}$$

and  $\lambda \geq 2$ . When  $k, l \geq 1$

$$w \rightsquigarrow w' \xrightarrow{(2.31)} \bar{2}(\bar{1}\bar{3})1 \Longrightarrow \bar{1}\bar{2}1 \longrightarrow \bar{1}\bar{2}$$

and  $\lambda \geq 2$ .

**Case 3-viii'.** When  $k = l = 0$ ,

$$w \rightsquigarrow w' = \bar{3} \bar{2} 1 = \bar{1} \bar{3} \bar{2} = \bar{2} \bar{1} \bar{2} \implies \bar{1} \bar{2}$$

Thus  $w$  has  $\lambda = 1$  if and only if  $w = \bar{3} \bar{2} 1^x$  for some  $x \geq 1$ . When  $k = 0$  and  $l \geq 1$ ,

$$w \rightsquigarrow w' = \bar{3} \bar{2} (123)^l 1 = \bar{1} \bar{3} \bar{2} (123)^l \implies \bar{1} \bar{2} (123)^l \xrightarrow{(2.15)} \dots \xrightarrow{(2.15)} \bar{1} \bar{2}$$

and  $\lambda \geq 2$ . When  $k \geq 1$  and  $l = 0$ ,

$$w \rightsquigarrow w' = \bar{3} \bar{2} (\bar{1} \bar{3} \bar{2})^k 1 \implies \bar{1} \bar{2} (\bar{1} \bar{3} \bar{2})^k \xrightarrow{(2.29)} \dots \xrightarrow{(2.29)} \bar{1} \bar{2}$$

and  $\lambda \geq 2$ . When  $k, l \geq 1$

$$w \rightsquigarrow w' \xrightarrow{(2.31)} \bar{3} \bar{2} (\bar{1} \bar{3}) 1 \implies \bar{1} \bar{2} \bar{1} \bar{3} = \bar{2} \bar{3} \bar{3} \bar{2} \implies \bar{2} \bar{3} = \bar{1} \bar{2}.$$

and  $\lambda \geq 2$ .

**Case 3-ix'.** When  $k = l = 0$ ,

$$w \rightsquigarrow w' = \bar{3} \bar{2} 12 = \bar{2} \bar{3} \bar{3} \bar{2} \implies \bar{2} \bar{3}.$$

Thus  $w$  has  $\lambda = 1$  if and only if  $w = \bar{3} \bar{2} 1^x 2^y$  for some  $x, y \geq 1$ . When  $k = 0, l \geq 1$ ,

$$\begin{aligned} w \rightsquigarrow w' &= \bar{3} \bar{2} (123)^l 12 = \bar{1} \bar{2} \bar{3} \bar{2} (123)^l = \bar{2} \bar{1} \bar{1} \bar{2} (123)^l \implies \bar{2} \bar{3} (123)^l \\ &= \bar{1} \bar{2} (123)^l \xrightarrow{(2.15)} \dots \xrightarrow{(2.15)} \bar{1} \bar{2} \end{aligned}$$

and  $\lambda \geq 2$ . When  $k \geq 1, l = 0$ ,

$$w \rightsquigarrow w' = \bar{3} \bar{2} (\bar{1} \bar{3} \bar{2})^k 12 \implies \bar{1} \bar{2} (\bar{1} \bar{3} \bar{2})^k \xrightarrow{(2.29)} \dots \xrightarrow{(2.29)} \bar{1} \bar{2}$$

and  $\lambda \geq 2$ . When  $k, l \geq 1$

$$w \rightsquigarrow w' \xrightarrow{(2.31)} \bar{3} \bar{2} (\bar{1} \bar{3}) 12 \implies \bar{1} \bar{2} (\bar{1} \bar{3}) = \bar{1} \bar{1} \bar{2} \bar{2} \implies \bar{1} \bar{2}$$

and  $\lambda \geq 2$ .

Table 1 summarizes all the words with  $\lambda = 1$ .

Words  $A_x, \dots, D_{x,y,z,w}$  are defined in Table 1. We can see that any word with  $\lambda = 1$  and having one component has one of the forms;  $A_x, \dots, D_{x,y,z,w}$ .

□

**Lemma 2.14** *The leading terms of the Alexander polynomials of  $\mathcal{K}_n, A_x, B_{x,y}, C_{x,y,z}$  and  $D_{x,y,z,w}$  are the following:*

$$\begin{aligned} \mathcal{K}_n; & \quad \pm(1 - 4t - 6t^2 + 8t^3 - \dots) \quad \text{if } n \geq 2, \\ A_x; & \quad \pm(1 - 3t + \dots) \quad \text{if } x \geq 2, \\ B_{x,y}; & \quad \pm(1 - 3t + \dots) \quad \text{if } x, y \geq 3, \\ C_{x,y,z}; & \quad \pm(1 - 5t + \dots) \quad \text{if } x, y, z \geq 2, \\ C_{1,2,z}, C_{1,y,2}, C_{2,y,1}, C_{x,2,1}; & \quad \pm(1 - 4t + 6t^2 - 7t^3 + \dots) \quad \text{if } x, y, z \geq 4, \\ C_{1,y,z}, C_{x,y,1}; & \quad \pm(1 - 4t + 7t^2 + \dots) \quad \text{if } x, y, z \geq 3 \\ D_{x,y,z,w}, D_{x,y,z,1}; & \quad \pm(1 - 6t + \dots) \quad \text{if } x, y, z, w \geq 2, \\ D_{x,y,1,w}; & \quad \pm(1 - 5t + \dots) \quad \text{if } x, y, w \geq 2. \end{aligned}$$

In particular,  $\mathcal{K}_n \neq A_x, B_{x,y}, C_{x,y,z}, D_{x,y,z,w}$ .

Table 1:

case	word with $\lambda = 1$ .
i	none.
ii	$\bar{2} \bar{1} \bar{3} 1^x$ (2 or 3 components.)
v	$\bar{3} 1^x 2^y 3^z 1^w = \begin{cases} C_{x+1,y,z} & \text{when } w = 1, \\ D_{x+1,y,z,w-1} & \text{when } w \geq 2. \end{cases}$ $\bar{3} \bar{3} 1^x$ (2 or 3 components.)
vi	$\bar{3} \bar{3} 1^x 2^y =: B_{x,y}$ . $\bar{3} 1^x 2^y 3^z 1^w 2^v = \begin{cases} C_{x+v+1,y,z} & \text{when } w = 1, \\ D_{x+v+1,y,z,w-1} & \text{when } w \geq 2. \end{cases}$
vii	$\bar{1} \bar{3} 1^x 2^y 3^z = \bar{1} \bar{3} 1^{x+z} 2^y$
ix	$\bar{1} \bar{3} 1^x 2^y = \begin{cases} \bar{3} \bar{3} 1^{x+1} & \text{when } y = 1 \text{ (2 or 3 components)} \\ B_{x+1,y-1} & \text{when } y \geq 2 \end{cases}$
i'	none.
iii'	none.
iv'	$\bar{2} 1^x 2^y 3^z =: C_{x,y,z}$ .
v'	$\bar{2} 1^x 2^y 3^z 1^w =: D_{x,y,z,w}$ . $\bar{2} \bar{2} 1^x$ (2 or 3 components.)
viii'	$\bar{3} \bar{2} 1^x =: A_x$ .
ix'	$\bar{3} \bar{2} 1^x 2^y = \begin{cases} \bar{3} \bar{3} 2^{y+1} & \text{when } x = 1, \text{ (2 or 3 components)} \\ B_{x-1,y+1} & \text{when } x \geq 2. \end{cases}$

**Proof of Lemma 2.14.** We prove that the Alexander polynomial of  $C_{x,y,z}$  for  $x, y, z \geq 2$  is  $\pm(1 - 5t + \dots)$ . Recall that Xu's Bennequin surface is a minimal genus Seifert surface. Let  $F$  be the Bennequin surface of  $C_{x,y,z}$  and choose a basis

$$\{u^{(1)}, u^{(2)}, u_1^{(3)}, \dots, u_{x-1}^{(3)}, u_1^{(4)}, \dots, u_{y-1}^{(4)}, u_1^{(5)}, \dots, u_{z-1}^{(5)}\}$$

for  $H_1(F)$  as in Figure 10, where  $u^{(k)}$  ( $k = 1, \dots, 5$ ) corresponds to loop  $(k)$ .




Figure 10: The Bennequin surface  $F$  of  $C_{x,y,z} = \bar{2}1^x 2^y 3^z$  and a basis for  $H_1(F)$

With respect to the basis, let  $V_{x,y,z}$  denote the Seifert matrix for  $C_{x,y,z}$ .

$$V_{x,y,z} = \left[ \begin{array}{c|c|c|c} & 1 & & 1 \\ \hline 1 & & -1 & \\ \hline & -1 & 1 & \\ & & -1 & \ddots \\ & & & \ddots & 1 \\ & & & & -1 \\ \hline & & & -1 & 1 \\ & & & & -1 & \ddots \\ & & & & & \ddots & 1 \\ & & & & & & -1 \\ \hline & & & & & -1 & 1 \\ & & & & & & -1 & \ddots \\ & & & & & & & \ddots & 1 \\ & & & & & & & & -1 \end{array} \right]$$

The empty spaces contain only 0's. The 3rd (resp. 4th, 5th) diagonal block has size  $(x-1) \times (x-1)$  (resp.  $(y-1) \times (y-1)$ ,  $(z-1) \times (z-1)$ ). The Alexander polynomial satisfies:

$$\Delta_{x,y,z}(t) = \det(V_{x,y,z}^T - tV_{x,y,z})$$

$$= \det \left[ \begin{array}{c|c|c|c} & 1-t & & -t \\ \hline 1-t & & t & \\ \hline & -1 & -1+t & -t \\ & & 1 & \ddots & \ddots \\ & & & \ddots & \ddots & -t \\ & & & & 1 & -1+t \\ \hline 1 & & & -1+t & -t \\ & & & 1 & \ddots & \ddots \\ & & & & \ddots & \ddots & -t \\ & & & & & 1 & -1+t \\ \hline & 1 & & & & -1+t & -t \\ & & & & & 1 & \ddots & \ddots \\ & & & & & & \ddots & \ddots & -t \\ & & & & & & & 1 & -1+t \end{array} \right]$$

Expanding it in the  $(x+1)$ th column, we have;

$$\Delta_{x,y,z}(t) = (-1+t)\Delta_{x-1,y,z}(t)$$

$$-(-t) \det \begin{bmatrix} & 1-t & & & -t & & \\ 1-t & & t & & & & -t \\ & -1 & -1+t & -t & & & \\ & & 1 & \ddots & -t & & \\ & & & 1 & -1+t & -t & \\ & & & & & 1 & \\ 1 & & & & -1+t & -t & \\ & & & & 1 & \ddots & -t \\ & & & & & 1 & -1+t \\ & & 1 & & & & -1+t & -t \\ & & & & & 1 & \ddots & -t \\ & & & & & & 1 & -1+t \end{bmatrix}$$

$$= (-1+t)\Delta_{x-1,y,z}(t) + t\Delta_{x-2,y,z}(t).$$

If  $\Delta_{i,y,z}(t) = (-1)^i(\alpha_0 + \alpha_1 t + \alpha_2 t^2 + \dots)$  for  $i = x-1$  and  $x-2$ , then

$$\begin{aligned} \Delta_{x,y,z}(t) &= (-1+t)(-1)^{x-1}(\alpha_0 + \alpha_1 t + \alpha_2 t^2 + \dots) + t(-1)^{x-2}(\alpha_0 + \alpha_1 t + \alpha_2 t^2 + \dots) \\ &= (-1)^x(\alpha_0 + \alpha_1 t + \alpha_2 t^2 + \dots). \end{aligned}$$

In fact,  $\Delta_{x,y,z}(t) = (-1)^{x+y+z}(1 - 5t + \dots)$  for all  $x, y, z \in \{2, 3\}$ . By induction,  $\Delta_{x,y,z}(t) = (-1)^{x+y+z}(1 - 5t + \dots)$  for all  $x, y, z \geq 2$ .

Other cases follow by similar arguments. □

Finally we are ready to prove the theorem.

**Proof of Theorem 2.8.** By Lemmas 2.10, 2.12, 2.14, our knot  $\mathcal{K}_{2m}$  where  $(m \geq 1)$  cannot be a 3-braid. Thus by Lemma 2.9, Theorem 2.8 follows. □

### 3 Uniqueness of the algebraic crossing number at minimal braid index

#### 3.1 Sharpness of the MFW inequality and conjectures

It has been conjectured (see [9] p.357 for example) that the exponent sum in a minimal braid representation is a knot invariant.

**Conjecture 3.1 (Main Conjecture):** *Let  $\mathcal{K}$  be a knot type of braid index  $b_{\mathcal{K}}$ . If  $K^1$  and  $K^2$  are braid representatives of  $\mathcal{K}$  with  $b_{K^1} = b_{K^2} = b_{\mathcal{K}}$  then their algebraic crossing numbers have  $c_{K^1} = c_{K^2}$*

We deform it into a stronger form:

**Conjecture 3.2** *Let  $\mathcal{B}_{\mathcal{K}}$  be the set of braid representatives of  $\mathcal{K}$ . Let  $\Phi : \mathcal{B}_{\mathcal{K}} \rightarrow \mathbb{N} \times \mathbb{Z}$  be a map such that  $\Phi(K) := (b_K, c_K)$  for  $K \in \mathcal{B}_{\mathcal{K}}$ . Then there exists a unique  $c_{\mathcal{K}} \in \mathbb{Z}$  with*

$$\Phi(\mathcal{B}_{\mathcal{K}}) = \{(b_{\mathcal{K}} + x + y, c_{\mathcal{K}} + x - y) \mid x, y \in \mathbb{N}\}, \tag{3.1}$$

the infinite quadrant region in Figure 11.




Figure 11: The region of braid representatives of  $\mathcal{K}$

The inclusion “ $\supset$ ” is trivial by the following argument: Let  $K_\star \in \mathcal{B}_\mathcal{K}$  be a minimal braid representative with  $\Phi(K_\star) = (b_\mathcal{K}, c_\mathcal{K})$ . Suppose  $K \in \mathcal{B}_\mathcal{K}$  is obtained from  $K_\star$  after applying (+)-stabilization  $x$ -times and then (-)-stabilization  $y$ -times. Then  $(b_\mathcal{K} + x + y, c_\mathcal{K} + x - y) = \Phi(K) \in \Phi(\mathcal{B}_\mathcal{K})$ .

The MFW inequality (1.2) says that  $c_K \geq -b_K + (d_+ + 1)$ ,  $c_K \leq b_K + (d_- - 1)$  for any  $K \in \mathcal{B}_\mathcal{K}$ . Thus

$$\Phi(\mathcal{B}_\mathcal{K}) \subset \{(x, y) \mid b_\mathcal{K} \leq x, -x + (d_+ + 1) \leq y \leq x + (d_- - 1)\}. \quad (3.2)$$

Before we provide examples of the conjectures we present:

**Theorem 3.3** *Sharpness of the MFW inequality implies the truth of Conjectures 3.1 and 3.2. In particular;*

$$b_\mathcal{K} = \frac{d_+ - d_-}{2} - 1, \quad c_\mathcal{K} = \frac{d_+ + d_-}{2}.$$

We remark that the statement in the theorem with regard to Conjecture 3.1 has been well known to many people.

**Proof of Theorem 3.3.** Let  $K_\star \in \mathcal{B}_\mathcal{K}$  be a minimal braid representative. Since the MFW inequality (1.2) is sharp on  $\mathcal{K}$ , we have  $c_{K_\star} - b_\mathcal{K} + 1 = d_-$ , and  $d_+ = b_\mathcal{K} + c_{K_\star} - 1$ , i.e.,  $c_{K_\star} = (d_+ + d_-)/2$  which is independent of the choice of  $K_\star$ . Thus we denote  $c_{K_\star} =: c_\mathcal{K}$ . In this case, the right side of (3.2) coincides with the right side of (3.1) and we have the other inclusion “ $\subset$ ” of (3.1).  $\square$

**Example 3.4** Both of the conjectures are true for unlinks, torus links, closed positive braids with a full twist (for example, the Lorenz links) [7], 2-bridge links and alternating fibered links [14], where the MFW inequality is sharp and one can apply Theorem 3.3.

Also Conjecture 3.1 applies to links with braid index  $\leq 3$  [3]. However, this case has been settled by a completely different way, the classification of 3-braids. Namely, any link of braid index 3 admits a unique conjugacy class of 3-braid representatives or has at most two conjugacy classes of 3-braid representatives related to each other by a flype move, which does not change the algebraic crossing number of the link.

Every transversal knot  $TK$  in  $S^3$  with the standard contact structure is transversally isotopic to a transversal closed braid  $K$  [1]. The *Bennequin number*  $\beta$  is an invariant of transversal knots. By

the identification of  $TK$  and  $K$ , we have  $\beta(K) = c_K - b_K$ . If Conjecture 3.2 is true for  $\mathcal{K}$ , then the *maximal* Bennequin number  $\beta_{\max}(\mathcal{K})$  for the knot type  $\mathcal{K}$  is realized on  $\mathcal{B}_{\mathcal{K}} \ni K$ 's with plotted vertices  $\Phi(K)$  on the upper half boundary of the quadrant region of Figure 11. Let  $\gamma(K) := c_K + b_K$ . For any  $K$  and its mirror image  $\overline{K}$ , we have  $\gamma(K) = -\beta(\overline{K})$ . Thus  $\gamma_{\min}(\mathcal{K}) = -\beta_{\max}(\overline{\mathcal{K}})$ . See Figure 11. Thus, investigation of  $\beta_{\max}(\mathcal{K})$  is related to Conjecture 3.2.

### 3.2 Cabling and the conjectures

In this subsection, we study behavior of the deficit of the MFW inequality under cabling and prove Theorem 3.5. As a consequence, we observe that the Conjecture 3.1 is true for many of the knots and links that appeared in Section 2, where the MFW-inequality is not sharp, i.e., we cannot apply Theorem 3.3.

We also prove, in Theorems 3.10 and 3.12, that the truth of Conjecture 3.1 is “inherited” through cabling operations.

Let us fix some notation. Let  $\mathcal{K}$  be a knot type. Denote the  $(p, q)$ -cable of  $\mathcal{K}$  by  $\mathcal{K}_{p,q}$ . Let  $K$  be a braid representative with  $b_K = b_{\mathcal{K}}$  and with algebraic crossing number  $c_K$ . Put

$$k := q - p \cdot c_K$$

and let  $K_{p,q}$  denote the  $p$ -parallel copies of  $K$  with a  $k/p$ -twist (see Figure 12). We can assume




Figure 12:  $(4, q)$ -cable ( $q = 4 \cdot 3 + k$ ) of the right hand trefoil

that  $K_{p,q}$  is on the boundary of a tubular neighborhood  $N$  of  $K$  (thus  $K$  is the core of solid torus  $N$ ). Then

$$q = \text{lk}(K_{p,q}, K)$$

where ‘lk’ is the linking number. Figure 12 shows that the algebraic crossing number of  $K_{p,q}$  is

$$c_{K_{p,q}} = p^2 c_K + k(p - 1) = q(p - 1) + p \cdot c_K. \quad (3.3)$$

Thanks to [20], we know that the braid index of  $\mathcal{K}_{p,q}$  satisfies

$$b_{\mathcal{K}_{p,q}} = p \cdot b_{\mathcal{K}}. \quad (3.4)$$

Although one can see a similar result in Theorem 7 of [18], we state the following for completeness of our discussion:

**Theorem 3.5** Suppose  $K^1$  and  $K^2$  are braid representatives of  $\mathcal{K}$  with  $b_{K^1} = b_{K^2} = b_{\mathcal{K}}$  and with distinct algebraic crossing numbers  $c_{K^1} < c_{K^2}$ , (i.e., Conjecture 3.1 does not apply to  $\mathcal{K}$ ). Then the deficit  $D_{\mathcal{K}_{p,q}}$  of the MFW inequality for  $(p, q)$ -cable  $\mathcal{K}_{p,q}$  is;

$$D_{\mathcal{K}_{p,q}} \geq \frac{p}{2}(c_{K^2} - c_{K^1}) \geq p. \quad (3.5)$$

**Proof of Theorem 3.5.** Thanks to (3.4), and by the construction of  $(K^1)_{p,q}$  and  $(K^2)_{p,q}$ , they are both minimal braid representatives of  $\mathcal{K}_{p,q}$  i.e.,  $b_{K^1_{p,q}} = b_{K^2_{p,q}} = b_{\mathcal{K}_{p,q}} = p \cdot b_{\mathcal{K}}$ .

Let  $k_1, k_2$  be integers satisfying  $q = pc_{K^1} + k_1 = pc_{K^2} + k_2$ . By (3.3) we have  $c_{K^1_{p,q}} = p^2c_{K^1} + k_1(p-1)$  and  $c_{K^2_{p,q}} = p^2c_{K^2} + k_2(p-1)$ . Therefore,  $c_{K^2_{p,q}} - c_{K^1_{p,q}} = p(c_{K^2} - c_{K^1})$ . By (1.2) we have  $c_{K^2_{p,q}} - b_{\mathcal{K}_{p,q}} + 1 \leq d_- \leq d_+ \leq c_{K^1_{p,q}} + b_{\mathcal{K}_{p,q}} - 1$ , and by Definition 2.1,

$$D_{\mathcal{K}_{p,q}} \geq \frac{1}{2}(c_{K^2_{p,q}} - c_{K^1_{p,q}}) = \frac{p}{2}(c_{K^2} - c_{K^1}). \quad (3.6)$$

This is the first inequality of (3.5).

Notice that  $K^1$  and  $K^2$  are related each other by a sequence of Markov moves [2]. Let  $K^1 = B_1 \rightarrow B_2 \rightarrow \dots \rightarrow B_n = K^2$  be a Markov tower. Each arrow corresponds to either braid isotopy, stabilization or destabilization moves. Let  $(x_i, y_i)$  be the braid index and the algebraic crossing number of  $B_i$ . Then  $(x_{i+1}, y_{i+1}) - (x_i, y_i) = (0, 0), (\pm 1, \pm 1)$  or  $(\mp 1, \pm 1)$  depending on the move corresponding to the arrow between  $B_{i+1}$  and  $B_i$ . Since  $x_1 = x_n = b_{\mathcal{K}}$  the difference  $c_{K^1} - c_{K^2} = y_1 - y_n \neq 0$  must be an even integer. Therefore, we get the second inequality of (3.5).  $\square$

**Corollary 3.6** Conjecture 3.1 is true for all  $9_{42}, 9_{49}, 10_{132}, 10_{150}, 10_{156}$ .

We remark that this follows also from Corollary 2 of [18].

**Proof.** Knotscape computes that the deficit of 2-cable  $\mathcal{K}_{2,2c_{\mathcal{K}}+1}$  is 1 for each knot.

$\mathcal{K}$	$b_{\mathcal{K}}$	$D_{\mathcal{K}}$	$K$	$c_K$	$D_{\mathcal{K}_{2,2c_{\mathcal{K}}+1}}$
$9_{42}$	4	1	$aaacBAAcB$	1	1
$9_{49}$	4	1	$aabbcbAbbcB$	7	1
$10_{132}$	4	2	$AbcaaaBBBcb$	3	1
$10_{150}$	4	1	$aabbcbABccB$	5	1
$10_{156}$	4	1	$aaacBAAcAb$	3	1

Comparing with (3.5), each  $\mathcal{K}$  must have unique algebraic crossing number.  $\square$

Thanks to Knotscape, the  $(2, 2c_{\mathcal{K}_n} + 1)$ -cable of  $\mathcal{K}_n = BM_{-1, -2, n, 2}$  has deficit = 1 if  $|n|$  is small. i.e., Conjecture 3.1 is true for  $\mathcal{K}_n$  if  $|n|$  is small. Corollary 3.6 implies:

**Corollary 3.7** Conjecture 3.1 is true for the prime links  $\mathcal{A}^n(9_{42})$  (see Figure 3).

**Proof.** We know that  $9_{42}$  has unique algebraic crossing number = 1 by Corollary 3.6. Since each link component of  $\mathcal{A}^n(9_{42})$  is  $9_{42}$ , we get this corollary.  $\square$

With regard to the deficit of cabled links, we conjecture that:

**Conjecture 3.8** For any  $q$ , the limit  $\lim_{p \rightarrow \infty} D_{\mathcal{K}_{p,q}}$  of deficits exists.

**Remark 3.9** If Conjecture 3.8 is true, then (3.5) of Theorem 3.5 implies the truth of Conjecture 3.1.

We present another property of cabling:

**Theorem 3.10** Let  $\mathcal{K}$  be a non-trivial knot type. If Conjecture 3.1 is true for  $\mathcal{K}$  then it is also true for  $\mathcal{K}_{p,q}$  when  $p \geq 2$ .

In particular, if  $c_{\mathcal{K}}$  and  $c_{\mathcal{K}_{p,q}}$  denote the unique algebraic crossing numbers of  $\mathcal{K}$  and  $\mathcal{K}_{p,q}$  respectively in their minimal braid representatives then we have

$$c_{\mathcal{K}_{p,q}} = (p-1)q + p \cdot c_{\mathcal{K}}.$$

**Remark 3.11** Suppose  $\mathcal{K}$  is the right hand trefoil. The MFW inequality is sharp on  $\mathcal{K}$ . Since its cable  $\mathcal{K}_{2,7}$  has deficit  $D_{\mathcal{K}_{2,7}} = 1$  (see [13]), we cannot apply Theorem 3.3. However Theorem 3.10 guarantees the truth of the conjecture for  $\mathcal{K}_{2,7}$ .

The following proof is inspired by the work of Williams [20], whose main result can be seen in formula (3.4). Note that his result holds not only for cable knots but also for generalized cable links. For the sake of completeness we repeat part of his discussion.

**Proof of Theorem 3.10.** Assume Conjecture 3.1 is true for  $\mathcal{K}$  and denote the unique algebraic crossing number at minimal braid index by  $c_{\mathcal{K}}$ .

Let  $K$  be a braid representative of  $\mathcal{K}$ . Suppose  $K'$  is a braid representative of  $\mathcal{K}_{p,q}$  on the boundary of a small tubular (solid torus) neighborhood  $N$  of  $K$ . We may regard the  $z$ -axis as the braid axis. Let  $\phi : \mathbb{R}^3 \rightarrow \mathbb{R}^3$  be a diffeomorphism of compact support so that  $(\star)$   $\phi(K') \subset \partial\phi(N)$  has exactly  $p \cdot b_{\mathcal{K}}$  maxima and  $p \cdot b_{\mathcal{K}}$  minima (both non-degenerate critical points) and no other critical points and  $(\star\star)$  the “height” function  $h : \partial\phi(N) \simeq T^2 \rightarrow \mathbb{R}$  is a Morse function. In particular,  $\phi(K')$  has a braid position with braid index  $p \cdot b_{\mathcal{K}}$ .

By  $(\star\star)$ , a generic intersection of the horizontal plane with  $\partial\phi(N) \simeq T^2$  consists of disjoint simple closed curves. Furthermore, these simple closed curves are either meridians of  $T^2$  or trivial in  $T^2$  since  $\mathcal{K}$  is knotted (Remark 1 of [20]).

Remark 2 in [20] says that there is a plane  $\pi$  (parallel to the  $(xz)$ -plane) intersecting transversely with  $T^2$  in a meridian.

Let  $J$  be an innermost one among such meridians. Then  $J$  bounds a disk  $d \subset \pi \cap \phi(N)$  which separates  $\phi(K')$  into arcs  $\{C_i\}$ . Close each  $C_i$  with aid of some arc  $D_i \subset d$  and set  $\hat{K}_i := C_i \cup D_i$ . See Figure 13.

Thanks to Remark 3 in [20],  $p$  of  $\hat{K}_i$ 's are non-trivial (i.e., do not bound any disk in  $\phi(N)$ ) since the linking number of  $J$  and  $\phi(K')$  pushed a little bit into the interior of  $\phi(N)$  is  $p$ .

Discard trivial  $\hat{K}_i$ 's.

Our  $\hat{K}_i$ 's are not in a braid position. As in [20], we make them have a braid position: Choose another plane  $\pi'$  just below  $\pi$  and call the annulus between the two planes  $A$  (see Figure 13). We may assume that the other boundary curve  $J' \subset \partial A$  is parallel to  $J$ .




Figure 13: Construction of  $K_i$  from  $\hat{K}_i$

As in the passage of Figure 13 replace the arc  $D_i \cup (C_i \cap A)$  (the left sketch) with  $D'_i \subset A$  (the right sketch) and construct  $p$ -parallels;

$$K_i := (C_i - (C_i \cap A)) \cup D'_i \subset \partial\phi(N) \quad \text{for } i = 1, \dots, p, \quad (3.7)$$

which is in a braid position. Also the  $K_i$ 's are disjoint from each other and each is isotopic to the core of the solid torus  $\phi(K) \simeq \mathcal{K}$ , thus  $b_{\mathcal{K}} \leq b_{K_i}$ . Then we have

$$\begin{aligned} p \cdot b_{\mathcal{K}} &\leq \sum_{i=1}^p \{b_{K_i} = \text{number of max of } K_i\} \\ &\leq \{\text{number of max of } \phi(K')\} = p \cdot b_{\mathcal{K}} \end{aligned}$$

where the last equality holds by  $(\star\star)$  above. This implies that  $(\dagger)$  there are no trivial  $\hat{K}_i$ 's (we didn't have to discard anything),  $(\dagger\dagger)$  each knot has  $b_{\mathcal{K}} = b_{K_i}$ . Let  $n, 0 \leq m < p$  be integers such that

$$q = p(c_{\mathcal{K}} + n) + m.$$

By  $(\dagger)$ , the  $p$ -component link  $L := K_1 \cup \dots \cup K_p$  is obtained from  $\mathbf{K}' := \phi(K')$  by using the meridian disk  $d$  to create a cutout and adding an  $m/p$ -twist along the annulus  $A$ , then gluing the end-points. See Figure 14. In other words,  $L$  is the  $(p, p(c_{\mathcal{K}} + n))$ -cable of  $K$ . From  $(\dagger\dagger)$  we have  $c_{K_i} = c_{\mathcal{K}} = c_{\mathcal{K}}$ .




Figure 14: From  $A \cap \mathbf{K}'$  to  $A \cap L$ , where  $p = 7, m = 2$

Therefore,  $L$  has the algebraic crossing number

$$\begin{aligned} c_L &= \sum_{i=1}^p c_{K_i} + \sum_{i \neq j} \text{lk}(K_i, K_j) = p \cdot c_{\mathcal{K}} + p(p-1)(c_{\mathcal{K}} + n) \\ &= p^2 c_{\mathcal{K}} + p(p-1)n \end{aligned}$$

and  $c_{\mathbf{K}'} - c_L = m(p - 1)$ . Thus,

$$\begin{aligned}
 c_{\mathbf{K}'} &= p^2 c_{\mathcal{K}} + p(p - 1)n + m(p - 1) \\
 &= p^2 c_{\mathcal{K}} + (p - 1)(pn + m) \\
 &= p^2 c_{\mathcal{K}} + (q - p \cdot c_{\mathcal{K}}) \\
 &= (p - 1)q + p \cdot c_{\mathcal{K}},
 \end{aligned}$$

which is independent of the choice of  $\mathbf{K}' \in \mathcal{B}_{\mathcal{K}_{p,q}}$ . Compare with (3.3). This concludes the uniqueness of the algebraic crossing number of  $\mathcal{K}_{p,q}$  at minimal braid index. □

A similar result to Theorem 3.10 holds for links.

**Theorem 3.12** *Let  $\mathcal{L} = \mathcal{K}^{(1)} \cup \dots \cup \mathcal{K}^{(l)}$  be an  $l$ -component link of braid index  $= b_{\mathcal{L}}$ . Assume that each  $\mathcal{K}^{(j)}$  is a non-trivial knot. Let  $\mathcal{L}' := \mathcal{K}_{p,q_1}^{(1)} \cup \dots \cup \mathcal{K}_{p,q_l}^{(l)}$  be the  $p$ -cable of  $\mathcal{L}$  such that  $q_j = \text{lk}(\mathcal{K}^{(j)}, \mathcal{K}_{p,q_j}^{(j)})$  for  $j = 1, \dots, l$ .*

*If  $\mathcal{L}$  and every component  $\mathcal{K}^{(j)}$  have unique algebraic crossing numbers  $c_{\mathcal{L}}, c_{\mathcal{K}^{(j)}}$  in minimal braid representations, then so does  $\mathcal{L}'$ .*

*Furthermore, let  $k_j$  satisfy  $q_j = p \cdot c_{\mathcal{K}^{(j)}} + k_j$  then*

$$c_{\mathcal{L}'} = p^2 c_{\mathcal{L}} + (p - 1)(k_1 + \dots + k_l). \quad (3.8)$$

**Remark 3.13** The assumption for  $\mathcal{K}^{(j)}$  in the second paragraph of Theorem 3.12 is essential by the following reason: If  $b_{\mathcal{L}} = \sum_{j=1}^l b_{\mathcal{K}^{(j)}}$ , then the existence of unique algebraic crossing number of  $\mathcal{L}$  in minimal braid representation implies that each  $\mathcal{K}^{(j)}$  also has unique algebraic crossing number. However, this is not true in general. For instance, assume that  $\mathcal{L} = \mathcal{K}^{(1)} \cup \mathcal{K}^{(2)}$  is a 2-component link and has two braid representatives related to each other by a flype move as in Figure 15. The thick




Figure 15: A flype move

gray arcs are parallel braid strands of  $\mathcal{L}$ . Braidings occur inside the boxes  $P, Q, R$ . In particular, box  $R$  contains even number of half twists of  $\mathcal{K}^{(1)}$  (dashed arc) and  $\mathcal{K}^{(2)}$  (black arc). The flype move preserves the number of braid strands and the algebraic crossing number of the link, but it changes the algebraic crossing numbers of link components  $\mathcal{K}^{(1)}, \mathcal{K}^{(2)}$ . Namely, in the passage from the left sketch to the right sketch, the algebraic crossing number of  $\mathcal{K}^{(1)}$  decreases by 1 and the one for  $\mathcal{K}^{(2)}$  increases by 1. (This means that a flype move in general cannot be a composition of exchange moves.)

**Proof of Theorem 3.12.** Suppose  $L = K^{(1)} \cup \cdots \cup K^{(l)}$  is a minimal braid representative of  $\mathcal{L} = \mathcal{K}^{(1)} \cup \cdots \cup \mathcal{K}^{(l)}$ , i.e.,  $b_L = b_{\mathcal{L}}$ . Let  $c_{x,y} := 2 \cdot \text{lk}(\mathcal{K}^{(x)}, \mathcal{K}^{(y)})$  for  $x \neq y$ . Then

$$c_{\mathcal{L}} = c_L = \sum_{1 \leq x < y \leq l} c_{x,y} + \sum_{j=1}^l c_{\mathcal{K}^{(j)}}. \quad (3.9)$$

Let  $k_j, n_j, 0 \leq m_j < p$  be integers with

$$q_j = p \cdot c_{\mathcal{K}^{(j)}} + k_j = p(c_{\mathcal{K}^{(j)}} + n_j) + m_j \quad \text{for } j = 1, \dots, l. \quad (3.10)$$

Williams proved that the braid index of  $\mathcal{L}' = \mathcal{K}_{p,q_1}^{(1)} \cup \cdots \cup \mathcal{K}_{p,q_l}^{(l)}$  is  $p \cdot b_{\mathcal{L}}$  [20]. Let  $L'$  be a minimal braid representative of  $\mathcal{L}'$ . Let  $N_j$  be a tubular neighborhood of  $K^{(j)}$ . Let  $\phi$  be a compact support diffeo morphism of  $\mathbb{R}^3$  such that  $\phi(L') =: K'^{(1)} \cup \cdots \cup K'^{(l)} \subset \partial\phi(N_1 \cup \cdots \cup N_l)$  has a minimal braid position.

As we did in (3.7), for each  $j = 1, \dots, l$ , construct  $p$  parallels  $K_1^{(j)}, \dots, K_p^{(j)} \subset \partial\phi(N_j)$  from each  $K'^{(j)}$  by cutout by an (inner most) meridian disk  $d_j \subset \phi(N_j)$  and adding an  $m_j/p$ -twist along annulus  $A_j \subset \partial\phi(N_j)$  then gluing. Thus,

$$\text{lk}(K_i^{(j)}, K^{(h)}) = \begin{cases} c_{\mathcal{K}^{(j)}} + n_j & \text{when } j = h, \\ \text{lk}(K^{(j)}, K^{(h)}) = \frac{1}{2}c_{j,h} & \text{otherwise.} \end{cases} \quad (3.11)$$

Let

$$L_i := K_i^{(1)} \cup \cdots \cup K_i^{(l)} \quad \text{for } i = 1, \dots, p.$$

Thanks to [20] we know that  $L_i \simeq L$  and  $b_{L_i} = b_L = b_{\mathcal{L}}$ . By assumption of Theorem 3.12, it follows  $c_{L_i} = c_L = c_{\mathcal{L}}$ . The  $(p \cdot l)$ -component link  $L_1 \cup \cdots \cup L_p$  has the algebraic crossing number;

$$\begin{aligned} c_{L_1 \cup \cdots \cup L_p} &= \sum_{i=1}^p c_{L_i} + \sum_{x \neq y} \text{lk}(L_x, L_y) \\ &= p \cdot c_{\mathcal{L}} + \sum_{i=1}^p (p-1) \text{lk}(L_i, L) \\ &= p \cdot c_{\mathcal{L}} + \sum_{i=1}^p (p-1) \left\{ \sum_{j=1}^l \text{lk}(K_i^{(j)}, K^{(j)}) + \sum_{x \neq y} \text{lk}(K_i^{(x)}, K^{(y)}) \right\} \\ &\stackrel{(3.11)}{=} p \cdot c_{\mathcal{L}} + p(p-1) \left\{ \sum_{j=1}^l (c_{\mathcal{K}^{(j)}} + n_j) + \sum_{x < y} c_{x,y} \right\} \\ &\stackrel{(3.9)}{=} p \cdot c_{\mathcal{L}} + p(p-1) \left( c_{\mathcal{L}} + \sum_{j=1}^l n_j \right) \\ &= p^2 c_{\mathcal{L}} + p(p-1) \left( \sum_{j=1}^l n_j \right). \end{aligned}$$

Since only the difference between  $L_1 \cup \dots \cup L_p$  and  $L'$  occur on the annuli  $A_1, \dots, A_l$ , we have

$$\begin{aligned}
c_{L'} &= c_{L_1 \cup \dots \cup L_p} + (p-1) \sum_{j=1}^l m_j \\
&= p^2 c_{\mathcal{L}} + (p-1) \sum_{j=1}^l (pn_j + m_j) \\
&\stackrel{(3.10)}{=} p^2 c_{\mathcal{L}} + (p-1) \sum_{j=1}^l k_j,
\end{aligned}$$

which is independent of the choice of braid representative  $L'$ .  $\square$

With regard to Conjecture 3.2 we have:

**Theorem 3.14** *Let  $\mathcal{L} = \mathcal{K}^{(1)} \cup \dots \cup \mathcal{K}^{(l)}$  be an  $l$ -component link satisfying all the assumptions in Theorem 3.12. If Conjecture 3.2 is true for  $\mathcal{L}$  then it is also true for its  $p$ -cable  $\mathcal{L}' := \mathcal{K}_{p,q_1}^{(1)} \cup \dots \cup \mathcal{K}_{p,q_l}^{(l)}$ .*

**Proof of Theorem 3.14.** Let  $L$  (resp.  $L'$ ) be a braid representative of  $\mathcal{L}$  (resp.  $\mathcal{L}'$ ). Take tubular neighborhoods  $N = N_1 \cup \dots \cup N_l$  of  $L$  (each  $N_j$  is a solid torus) and let  $\phi : \mathbb{R}^3 \rightarrow \mathbb{R}^3$  be a compact support diffeo morphism such that  $\phi(L) =: K^{(1)} \cup \dots \cup K^{(l)}$ ,  $\phi(L') =: K'^{(1)} \cup \dots \cup K'^{(l)}$  have braid positions. They are not necessarily minimal braid representatives and in general  $b_{\phi(L')} \neq p \cdot b_{\phi(L)}$ . We may assume that  $K'^{(j)} \subset \partial\phi(N_j) \simeq T^2$ .

Let plane  $\pi_j = \{(x, y, z) | y = y_0\}$ , innermost meridian loop  $J_j \subset \pi_j \cap \partial\phi(N_j)$ , and meridian disk  $d_j \subset \pi_j \cap \phi(N_j)$  be as in the proof of Theorem 3.10. We may assume that the braid axis is *not* contained in  $\pi_j$  i.e.,  $y_0 \neq 0$ .

We deform  $K'^{(j)}$  in the following way: Suppose sub-arcs  $u \subset K'^{(j)}$  and  $v \subset J_j$  bound a disk  $\mathcal{D} \subset \partial\phi(N_j) \simeq T^2$ . If  $\mathcal{D}$  is innermost, then replace  $u$  with  $v$ . Repeat this until  $K'^{(j)}$  and  $J_j$  do not bound any disk in  $T^2$ . Add up all the linking numbers of  $\partial\mathcal{D}$ 's with the  $z$ -axis and denote it by  $x^{(j)} \geq 0$ .

Next, from the deformed  $K'^{(j)}$  above, construct  $p$ -parallels  $K_1'^{(j)}, \dots, K_p'^{(j)}$  as in (3.7). Since the plane  $\pi_j$  does not contain the  $z$ -axis, the  $m_j/p$ -twist along a thin annulus does not change the number of braid strands.

Suppose that  $b_{K^{(j)}} = b_{\mathcal{K}^{(j)}} + y^{(j)}$  and  $b_{K_i'^{(j)}} = b_{\mathcal{K}^{(j)}} + y^{(j)} + z_i^{(j)}$  with  $y^{(j)}$ ,  $y^{(j)} + z_i^{(j)} \geq 0$ . Let  $L_i := K_i^{(1)} \cup \dots \cup K_i^{(l)}$  then

$$b_{L_i} = \sum_{j=1}^l b_{K_i'^{(j)}} = b_{\mathcal{L}} + \sum_{j=1}^l (y^{(j)} + z_i^{(j)}), \quad (3.12)$$

$$b_{\phi(L')} = \sum_{i=1}^p b_{L_i} + \sum_{j=1}^l x^{(j)} \stackrel{(3.12)}{=} p \cdot b_{\mathcal{L}} + \sum_{j=1}^l (x^{(j)} + p \cdot y^{(j)} + \sum_{i=1}^p z_i^{(j)}). \quad (3.13)$$

Since  $L_i \simeq \mathcal{L}$ , our assumption of this theorem and (3.12) give us

$$c_{\mathcal{L}} - \sum_{j=1}^l (y^{(j)} + z_i^{(j)}) \leq c_{L_i} \leq c_{\mathcal{L}} + \sum_{j=1}^l (y^{(j)} + z_i^{(j)}). \quad (3.14)$$

As in the proof of Theorem 3.12, let  $k_j, n_j, 0 \leq m_j < p$  satisfy  $q_j = p \cdot c_{\mathcal{K}^{(j)}} + k_j = p(c_{\mathcal{K}^{(j)}} + n_j) + m_j$ . Then we have

$$\begin{aligned}
c_{L_1 \cup \dots \cup L_p} &= \sum_{i=1}^p c_{L_i} + \sum_{x \neq y} \text{lk}(L_x, L_y) \\
&\stackrel{(3.14)}{\leq} p(c_{\mathcal{L}} + \sum_{j=1}^l y^{(j)}) + \sum_{i=1}^p \sum_{j=1}^l z_i^{(j)} + p(p-1)(c_{\mathcal{L}} + \sum_{j=1}^l n_j) \\
&= p^2 c_{\mathcal{L}} + p(p-1)(\sum_{j=1}^l n_j) + \sum_{j=1}^l (p \cdot y^{(j)} + \sum_{i=1}^p z_i^{(j)}),
\end{aligned}$$

and

$$\begin{aligned}
c_{\phi(L')} &\leq c_{L_1 \cup \dots \cup L_p} + \sum_{j=1}^l m_j + \sum_{j=1}^l x^{(j)} \\
&\leq p^2 c_{\mathcal{L}} + (p-1) \sum_{j=1}^l k_j + \sum_{j=1}^l (x^{(j)} + p \cdot y^{(j)} + \sum_{i=1}^p z_i^{(j)}). \tag{3.15}
\end{aligned}$$

Similarly,

$$p^2 c_{\mathcal{L}} + (p-1) \sum_{j=1}^l k_j - \sum_{j=1}^l (x^{(j)} + p \cdot y^{(j)} + \sum_{i=1}^p z_i^{(j)}) \leq c_{\phi(L')}. \tag{3.16}$$

We conclude the theorem by (3.8), (3.13), (3.15) and (3.16).  $\square$

**Corollary 3.15** *Conjectures 3.1 and 3.2 apply to iterated torus knots.*

**Proof of Corollary 3.15.** We know that the both conjectures apply to torus knots (Example 3.4). Thanks to Theorems 3.10, 3.12 and 3.14, we have this corollary.  $\square$

### 3.3 Connect sum and the conjecture

We will prove the following:

**Theorem 3.16** *If Conjecture 3.1 is true for knot types  $\mathcal{K}^1$  and  $\mathcal{K}^2$  then it is also true for the connect sum  $\mathcal{K}^1 \sharp \mathcal{K}^2$ .*

*In particular, denoting the unique algebraic crossing numbers of  $\mathcal{K}^i$  in minimal braid representatives by  $c_{\mathcal{K}^i}$  we have*

$$c_{\mathcal{K}^1 \sharp \mathcal{K}^2} = c_{\mathcal{K}^1} + c_{\mathcal{K}^2}.$$

Before we prove Theorem 3.16 let us recall two important known results:

**Lemma 3.17** ([10] Theorem 2.12.) *Up to ordering of summands, there is a unique expression for a knot type  $\mathcal{K}$  as a finite connect sum of prime knots.*

**Lemma 3.18 (The composite braid theorem, [4].)** *Let  $\mathcal{K}$  be a composite link, and let  $K$  be an arbitrary closed  $n$ -braid representative of  $\mathcal{K}$ . Then there is an obvious composite  $n$ -braid representative  $K^\bullet$  of  $\mathcal{K}$  (see Figure 16) and a finite sequence of closed  $n$ -braids:*

$$K = K_0 \rightarrow K_1 \rightarrow \cdots \rightarrow K_m = K^\bullet$$

*such that  $K_{i+1}$  is obtained from  $K_i$  by either braid isotopy or an exchange move.*




Figure 16: An obvious composite braid

**Proof of Theorem 3.16.** Since an exchange move does not change the algebraic crossing number, Lemmas 3.17 and 3.18 imply the truth of Theorem 3.16.  $\square$

As a corollary of Theorem 3.16 we have:

**Theorem 3.19** *If Conjecture 3.2 is true for  $\mathcal{K}^1, \mathcal{K}^2$  then it is also true for  $\mathcal{K}^1 \# \mathcal{K}^2$ .*

**Proof of Theorem 3.19.** Let  $K$  be a braid representative of  $\mathcal{K}^1 \# \mathcal{K}^2$ . By Lemma 3.18, after applying exchange moves and braid isotopy to  $K$  one can get a composite braid representative  $K^\bullet = K^1 \# K^2$ . Suppose  $b_{K^i} = b_{\mathcal{K}^i} + x_i$  with  $x_i \geq 0$ . Then

$$\begin{aligned} b_K &= b_{K^\bullet} = b_{K^1} + b_{K^2} - 1 = (b_{\mathcal{K}^1} + b_{\mathcal{K}^2} - 1) + (x_1 + x_2) \\ &= b_{\mathcal{K}^1 \# \mathcal{K}^2} + (x_1 + x_2). \end{aligned} \tag{3.17}$$

Our assumption gives  $c_{\mathcal{K}^i} - x_i \leq c_{K^i} \leq c_{\mathcal{K}^i} + x_i$ . Since  $c_K = c_{K^\bullet} = c_{K^1} + c_{K^2}$ , we have  $(c_{\mathcal{K}^1} + c_{\mathcal{K}^2}) - (x_1 + x_2) \leq c_K \leq (c_{\mathcal{K}^1} + c_{\mathcal{K}^2}) + (x_1 + x_2)$ . Thanks to Theorem 3.16,

$$c_{\mathcal{K}^1 \# \mathcal{K}^2} - (x_1 + x_2) \leq c_K \leq c_{\mathcal{K}^1 \# \mathcal{K}^2} + (x_1 + x_2). \tag{3.18}$$

The truth of Conjecture 3.2 follows by (3.17), (3.18).  $\square$

## References

- [1] Bennequin, D., *Entrelacements et équations de Pfaff* *Astrisque*, **107-108**, (1983) 87–161.
- [2] Birman, J. S., *Braids, links, and mapping class groups*, *Annals of Mathematics Studies*, No. 82. Princeton University Press, Princeton, N.J.; University of Tokyo Press, Tokyo, 1974.
- [3] Birman, J. S. and Menasco, W. W., *Studying links via closed braids III: Classifying links which are closed 3-braids*, *Pacific J. Math.* **161**, (1993), no. 1, 25–113.

- [4] Birman, J. S. and Menasco, W. W., *Studying links via closed braids. IV. Composite links and split links*, Invent. Math. 102 (1990), no. 1, 115–139 Erratum: Invent. Math. 160 (2005), no. 2, 447–452.
- [5] Birman, J. S. and Menasco, W. W., *Studying links via closed braids. V. The unlink*, Trans. Amer. Math. Soc. **329** (1992), no. 2, 585–606.
- [6] Birman, J. S. and Menasco, W. W., *Stabilization in the braid groups I: MTWS*, preprint.
- [7] Franks, J. and Williams, R. F., *Braids and the Jones Polynomial*, Trans. Amer. Math. Soc., **303**, (1987), 97–108.
- [8] Giroux, E., *Gomtrie de contact: de la dimension trois vers les dimensions suprieures*. Proceedings of the International Congress of Mathematicians, Vol. II (Beijing, 2002), 405–414, Higher Ed. Press, Beijing, 2002.
- [9] Jones, V. F. R., *Hecke algebra representations of braid groups and link polynomials*, Ann. of Math., **126**, (1987), 335–388.
- [10] Lickorish, W. B., *An introduction to knot theory*, Graduate Texts in Mathematics, **175**, Springer-Verlag, New York, 1997.
- [11] Melvin, P. M. and Morton, H. R., *Fibred knots of genus 2 formed by plumbing Hopf bands* Proc. London Math. Soc. (2) **34** (1986), 159–168.
- [12] Morton, H. R., *Seifert circles and knot polynomials*, Math. Proc. Cambridge Philos. Soc., **99**, (1986), 107–109.
- [13] Morton, H. R. and Short, H. B., *The 2-variable polynomial of cable knots* Math. Proc. Cambridge Philos. Soc. 101 (1987), no. 2, 267–278.
- [14] Murasugi, K., *On the braid index of alternating links* Trans. Amer. Math. Soc. **326** (1991), no. 1, 237–260.
- [15] Neumann, W. D., Private communication.
- [16] Neumann, W. D. and Rudolph, L., *Difference index of vector fields and the enhanced Milnor number* Topology **29** (1990), no. 1, 83–100.
- [17] Neumann, W. D. and Zagier, D. *Volumes of hyperbolic three-manifolds* Topology **24** (1985), no. 3, 307–332.
- [18] Stoimenow, A., *On the crossing number of positive knots and braids and braid index criteria of Jones and Morton-Williams-Franks* Trans. Amer. Math. Soc. 354 (2002), no. 10, 3927–3954
- [19] Stoimenow, A., *Properties of closed 3-braids* preprint.
- [20] Williams, R. F., *The braid index of generalized cables*, Pacific J. Math. **155** (1992), no. 2, 369–375.
- [21] Xu, Peijun, *The genus of closed 3-braids*, J. Knot Theory Ramifications **1** (1992), no. 3, 303–326.

Department of Mathematics, Columbia University, New York, NY 10027  
*E-mail address:* yuri@math.columbia.edu